

Crossed Andreev reflection and charge imbalance in diffusive NSN structures

Dmitri S. Golubev¹, Mikhail S. Kalenkov², and Andrei D. Zaikin^{1,2}

¹*Forschungszentrum Karlsruhe, Institut für Nanotechnologie, 76021 Karlsruhe and*

²*I.E. Tamm Department of Theoretical Physics, P.N. Lebedev Physics Institute, 119991 Moscow, Russia*

We formulate a microscopic theory of non-local electron transport in three-terminal diffusive normal-superconducting-normal (NSN) structures with arbitrary interface transmissions. At low energies ε we predict strong enhancement of non-local spectral conductance $g_{12} \propto 1/\varepsilon$ due to quantum interference of electrons in disordered N-terminals. In contrast, non-local resistance R_{12} remains smooth at small ε and, furthermore, is found to depend neither on parameters of NS interfaces nor on those of N-terminals. At higher temperatures R_{12} exhibits a peak caused by the trade-off between charge imbalance and Andreev reflection. Our results are in a good agreement with recent experimental observations and can be used for quantitative analysis of future experiments.

In hybrid NS structures quasiparticle current flowing in a normal metal is inevitably converted into that of Cooper pairs inside a superconductor. For quasiparticle energies above the superconducting gap $\varepsilon > \Delta$ this conversion is accompanied by electron-hole (or charge) imbalance¹ which relaxes inside a superconductor at a typical inelastic length usually denoted as Λ_{Q^*} . As a result, at temperatures near the critical one T_C an electric field penetrates into a superconductor causing resistance enhancement for NS structures under consideration.

At subgap energies $\varepsilon < \Delta$ the physical picture becomes entirely different. In this case quasiparticle-to-Cooper-pair current conversion is provided by the mechanism of Andreev reflection²: A quasiparticle enters the superconductor from the normal metal at a length of order of the superconducting coherence length ξ_S , forms a Cooper pair together with another quasiparticle, while a hole goes back into the normal metal. Due to this process subgap conductance of the NS structure remains non-zero down to $T = 0$ ³. Furthermore, in the presence of disorder this subgap conductance can be greatly enhanced at low energies due to quantum interference effects^{4,5,6}.

Further interesting effects may occur in three-terminal NSN structures. Provided the distance between two N-terminals is smaller than or comparable with ξ_S , electrons penetrating into the superconductor from the first N-terminal may form Cooper pairs with electrons from the second N-terminal. Then a hole goes into the *second* N-metal making the charge transfer effectively non-local. This important phenomenon of non-local (or crossed) Andreev reflection (CAR)⁷ enables direct experimental realization of entanglement between electrons from spatially separated N-terminals.

CAR was detected and investigated in several recent experiments^{8,9,10,11} by measuring the non-local resistance of multiterminal NSN systems. The authors observed a rich structure of different features many of which are still waiting for their theoretical interpretation. Note that not only CAR but also other physical processes contribute to the non-local conductance g_{12} thus making this interpretation rather complicated. For instance, the contribution of elastic cotunneling (EC) to g_{12} exactly cancels that of CAR in the lowest order in NS interface transmissions




FIG. 1: NSN structure under consideration.

and at subgap energies¹². This cancellation is lifted either in higher orders in barrier transmissions¹³ or in the presence of interactions, e.g., with an effective external environment¹⁴, or under external ac bias¹⁵.

Another important issue is the effect of disorder in metallic terminals which needs to be analyzed for adequate interpretation of experimental results^{8,9,10,11}. Although CAR in disordered NSN structures was already addressed in a number of theoretical works^{16,17,18,19,20} in various physical limits, we believe that general analysis of this issue is still missing in the literature. For instance, the role of disorder-induced electron interference^{4,5,6} in the non-local subgap transport, the effect of high NS barrier transmissions on CAR as well as some other features remain unclear. Yet another important unresolved problem is to describe an interplay between CAR and non-local charge imbalance. It was demonstrated both experimentally^{8,10} and theoretically^{20,21} that such interplay may result in a large non-local resistance peak which occurs at temperatures slightly below the critical one. It was conjectured¹⁰ that the behavior of this peak is controlled by the charge imbalance length Λ_{Q^*} parametrically exceeding the length scale $\sim \xi_S$ relevant for CAR. In this work we develop a general theory of non-local electron transport in diffusive NSN structures which enables one to clarify the above issues and to formulate predictions to be tested in future experiments.

Model and basic equations. In what follows we will analyze a multiterminal diffusive NSN structure schematically shown in Fig. 1. Two normal terminals N_1 and N_2 with resistances r_{N_1} and r_{N_2} and electric poten-

tials V_1 and V_2 are connected to a superconducting electrode of length L with normal state (Drude) resistance r_L and electric potential $V = 0$ via small NS barriers with resistances R_1 and R_2 which can be expressed via channel transmissions $T_{1,n}$ and $T_{2,n}$ of these barriers as $1/R_{1(2)} = (e^2/\pi) \sum_n T_{1(2),n}$. For the sake of definiteness in Fig. 1 we chose specific geometry directly related, e.g., to experiments¹⁰ where the superconductor was fabricated in the form of a rather thin strip. NS barriers are located at the points $\mathbf{r}_{1,2} = (x_{1,2}, 0, 0)$ and the corresponding segments of a superconducting strip have normal state resistances r_{x_1} and r_{L-x_2} , see Fig. 1.

Our analysis is based on the quasiclassical Usadel equations for the Green-Keldysh matrix functions $\check{G}$. In the absence of interactions these equations read²²

$$iD\nabla(\check{G}\nabla\check{G}) = [\check{\Sigma}, \check{G}], \quad \check{G}^2 = 1, \quad (1)$$

where $[\check{a}, \check{b}] = \check{a}\check{b} - \check{b}\check{a}$, D is the diffusion constant and

$$\check{G} = \begin{pmatrix} \hat{G}^R & \hat{G}^K \\ 0 & \hat{G}^A \end{pmatrix}, \quad \check{\Sigma} = \check{1} \begin{pmatrix} \varepsilon + eV & \Delta \\ -\Delta^* & -\varepsilon + eV \end{pmatrix} \quad (2)$$

are 4×4 matrices in Keldysh $\otimes$ Nambu space, ε is the quasiparticle energy, $\Delta(T)$ is the superconducting order parameter which will be considered real further below and V is the electric potential.

Far from the interfaces between metals the quasiclassical Green functions $\check{G}$ coincide with their bulk equilibrium values. Deep in the superconductor they read

$$\hat{G}_S^{R,A} = \pm \frac{\hat{\tau}_3 \varepsilon + i\hat{\tau}_2 \Delta}{\sqrt{(\varepsilon \pm i\delta)^2 - \Delta^2}}, \quad \hat{G}_S^K = (\hat{G}_S^R - \hat{G}_S^A)n(\varepsilon), \quad (3)$$

where $n(\varepsilon) = \tanh(\varepsilon/2T)$ and $\hat{\tau}_i$ are Pauli matrices. In the normal terminals far from the tunnel barriers one has

$$\hat{G}_{1,2}^K = 2 \begin{pmatrix} \tanh \frac{\varepsilon + eV_{1,2}}{2T} & 0 \\ 0 & -\tanh \frac{\varepsilon - eV_{1,2}}{2T} \end{pmatrix}, \quad (4)$$

while the retarded and advanced Green functions $\hat{G}_{1,2}^{R,A}$ are set by the first Eq. (3) with $\Delta = 0$. In the vicinity of the barriers the Green functions deviate from the above equilibrium values and should be determined from Eqs. (1) supplemented by appropriate boundary conditions describing electron transfer across metallic interfaces. For diffusive superconductors one finds²³

$$\begin{aligned} \mathcal{A}_1 \sigma_1 \check{G}_1 \partial_x \check{G}_1 &= \mathcal{A}_1 \sigma_S \check{G}_S \partial_x \check{G}_S \\ &= \frac{e^2}{\pi} \sum_n \frac{2T_{1,n}[\check{G}_1, \check{G}_S]}{4 + T_{1,n}(\{\check{G}_1, \check{G}_S\} - 2)} \end{aligned} \quad (5)$$

for the first interface and similarly for the second one. Here $\mathcal{A}_{1,2}$ are the barrier cross sections and $\sigma_{S,1,2}$ are Drude conductivities of S- and N-terminals.

Having derived the Green-Keldysh functions $\check{G}$ one can easily evaluate the current density $\mathbf{j}$ in our system with the aid of the standard relation

$$\mathbf{j} = -\frac{\sigma}{8e} \int \text{tr}[\hat{\tau}_3(\check{G}\nabla\check{G})^K] d\varepsilon. \quad (6)$$

Non-local spectral conductance. The above general formalism enables one to describe electron transport at arbitrary barrier transmissions $T_{1,n}$ and $T_{2,n}$. Here we only assume that both NS barriers are sufficiently small to provide $R_{1,2} \gg r = \max(r_L, r_{N_1}, r_{N_2})$. This condition allows to effectively linearize Eqs. (1) and express the solution of linearized Usadel equations via the diffuson $\mathcal{D}^{rr'}(\omega)$ and the Cooperon $\mathcal{C}^{rr'}(\omega)$. The diffuson satisfies the following diffusion equation

$$\left(-i\omega + \frac{1}{\tau_{Q^*}} - D\nabla^2\right) \mathcal{D}^{rr'}(\omega) = \delta(\mathbf{r} - \mathbf{r}'), \quad (7)$$

while Cooperon is the solution of Eq. (7) with effective charge imbalance relaxation time τ_{Q^*} replaced by dephasing time τ_φ . At $T \sim T_C$ τ_{Q^*} depends on the electron inelastic relaxation time τ_{in} as¹ $\tau_{Q^*} \sim \tau_{in}T/\Delta(T)$.

Let us employ the standard representation of the Keldysh function $\hat{G}^K = \hat{G}^R \hat{h} - \hat{h} \hat{G}^A$ with $\hat{h} = f_L \hat{1} + f_T \hat{\tau}_3$, where f_L and f_T are respectively symmetric and anti-symmetric in energy parts of the distribution function. Combining the above expression for $\hat{G}^K$ with Eq. (6) we define the current across the first barrier

$$I_1 = \frac{1}{2e} \int d\varepsilon g_1(\varepsilon) [f_T^{N_1}(\varepsilon, \mathbf{r}_1) - f_T^S(\varepsilon, \mathbf{r}_1)]. \quad (8)$$

The spectral conductance $g_1(\varepsilon)$ is expressed via the functions $\hat{G}^R, \hat{G}^A$. Solving Eq. (1) for $\hat{G}^R$ and keeping terms up to the first order in $r/R_{1,2}$, we find

$$\begin{aligned} g_1(\varepsilon) &= g_1^{\text{BTK}}(\varepsilon) + \frac{\theta(\Delta - |\varepsilon|)\Delta^2 \text{Re} \mathcal{C}_1^{r_1 r_1}(2\varepsilon)}{\Omega^2 \frac{2e^2 N_1 R_1^2}{2e^2 N_1 R_1^2}} \\ &+ \frac{\Delta^2}{\Omega^2} \sum_{j=1,2} \frac{\text{Re} \mathcal{C}_S^{r_1 r_j}(2W(\varepsilon))}{2e^2 N_S R_j}, \end{aligned} \quad (9)$$

where $W(\varepsilon) = i\Omega = i\sqrt{\Delta^2 - \varepsilon^2}$ for $|\varepsilon| < \Delta$, $W(\varepsilon) = |\Omega| \text{sign } \varepsilon$ for $|\varepsilon| > \Delta$, and $g_1^{\text{BTK}}(\varepsilon)$ is defined by the standard expression³

$$\begin{aligned} g_1^{\text{BTK}}(\varepsilon) &= \frac{e^2}{\pi} \sum_n \left[\frac{2T_{1,n}^2 \theta(\Delta - |\varepsilon|)\Delta^2}{T_{1,n}^2 \varepsilon^2 + (2 - T_{1,n})^2 \Omega^2} \right. \\ &\quad \left. + \frac{2T_{1,n} \theta(|\varepsilon| - \Delta)|\varepsilon|}{T_{1,n}|\varepsilon| + (2 - T_{1,n})|\Omega|} \right]. \end{aligned} \quad (10)$$

Note that the terms $\propto \mathcal{C}_S, \mathcal{C}_1$ in Eq. (9) are evaluated in the limit $T_{1,n}, T_{2,n} \ll 1$ where they only matter as compared to $g_1^{\text{BTK}}(\varepsilon)$ provided $R_{1,2} \gg r$. The Cooperon term $\propto \mathcal{C}_1$ describes enhancement of Andreev conductance by electron interference in diffusive N-metal^{4,5,6}, while the term $\propto \mathcal{C}_S$ accounts for broadening of the density of states in the superconductor. The spectral conductance $g_2(\varepsilon)$ is given by Eq. (9) with interchanged indices $1 \leftrightarrow 2$.

Our next step is to solve the kinetic equation for the distribution function f_T . In the limit $r/R_{1,2} \rightarrow 0$ this solution is trivial: $f_T^S(\varepsilon) = 0$ and $f_T^{N_j}(\varepsilon, \mathbf{r}_j) = h(\varepsilon, V_j) \equiv$

$(\tanh[(\varepsilon + eV_j)/2T] - \tanh[(\varepsilon - eV_j)/2T])/2$ ($j = 1, 2$). In the first order in $r/R_{1,2}$ the function f_T^S is determined from the diffusion equation

$$(2\tilde{\Omega} - D\nabla^2)f_T = \sum_{j=1,2} \frac{g_j(\varepsilon)h(\varepsilon, V_j)}{2e^2N_S K(\varepsilon)} \delta(\mathbf{r} - \mathbf{r}_j), \quad (11)$$

where $K(\varepsilon) = \theta(\Delta - |\varepsilon|)\Delta^2/\Omega^2 - \theta(|\varepsilon| - \Delta)\varepsilon^2/\Omega^2$, and $\tilde{\Omega} = \theta(\Delta - |\varepsilon|)\Omega$. Resolving Eq. (11) and substituting the result into Eq. (8), we obtain

$$I_1(V_1, V_2) = \int d\varepsilon [g_{11}(\varepsilon)h(\varepsilon, V_1) - g_{12}(\varepsilon)h(\varepsilon, V_2)], \quad (12)$$

where

$$g_{11}(\varepsilon) = g_1(\varepsilon) - \frac{D_S^{r_1 r_1} (2i\tilde{\Omega})}{2e^2 N_S} \frac{g_1^2(\varepsilon)}{K(\varepsilon)} - \frac{D_1^{r_1 r_1} (0)}{2e^2 N_S} g_1^2(\varepsilon), \quad (13)$$

$$g_{12}(\varepsilon) = g_{21}(\varepsilon) = \frac{D_S^{r_1 r_2} (2i\tilde{\Omega})}{2e^2 N_S} \frac{g_1(\varepsilon)g_2(\varepsilon)}{K(\varepsilon)}, \quad (14)$$

and similarly for the current I_2 . The last two terms in the local conductance $g_{11}(\varepsilon)$ (13) describe partial conductance suppression respectively due to local charge imbalance inside the superconductor and due to non-equilibrium quasiparticles in the normal metal. At energies $|\varepsilon| > \Delta$ Eq. (14) accounts for the effect of *non-local* charge imbalance which yields non-zero contribution to $g_{12}(\varepsilon)$ already in the lowest order in $1/R_1 R_2$. In contrast, at subgap energies this lowest order contribution vanishes identically manifesting the well known cancellation between EC and CAR terms¹². This cancellation is lifted in higher orders in barrier transmissions¹³. Accordingly, the full expression for $g_{12}(\varepsilon)$ (14) does not vanish also for $|\varepsilon| < \Delta$ and describes non-trivial interplay between CAR and direct electron transfer in the presence of disorder.

Eq. (14) for the non-local spectral conductance – together with Eqs. (9), (10) and (13) – is the central result of this work. Note that this result is not specific to particular geometry of Fig. 1 but applies for other diffusive NSN structures as well.

Here, however, we will only analyze the system with effectively quasi-one-dimensional superconducting and normal wires, as shown in Fig. 1. Assuming $x_2 > x_1$ we obtain

$$\begin{aligned} D_S^{x_1 x_2}(\omega) &= \frac{\sinh[k(L - x_2)] \sinh kx_1}{kS_S D_S \sinh(kL)}, \\ C_j^{x_j x_j}(\omega) &= \frac{\tanh\left(\sqrt{(-i\omega + 1/\tau_\varphi)/D_j} L_j\right)}{S_j D_j \sqrt{(-i\omega + 1/\tau_\varphi)/D_j}}, \end{aligned} \quad (15)$$

where $j = 1, 2$. Here $S_{S,1,2}$ and $D_{S,1,2}$ are respectively effective cross sections and diffusion coefficients of the corresponding terminals and $k = \sqrt{(-i\omega + 1/\tau_{Q^*})/D_S}$. Substituting Eqs. (15) into (13)-(14) we arrive at the conductance matrix describing the system in Fig. 1.




FIG. 2: Non-local spectral conductance $g_{12}(\varepsilon)$ (normalized by $G_0 = r_{x_1} r_{L-x_2}/r_L R_1 R_2$) for diffusive NSN structures. We set $L = 10\sqrt{2}\xi_S(0)$, $x_1 = 0.45L$, and $x_2 = 0.55L$. (a) The case of two identical barriers with resistances $R_1 = R_2 = \pi/e^2 N_{ch} \tau$ (N_{ch} is the number of channels and τ is the barrier transmission) and for $r_{N_1} = r_{N_2} = 0$. (b) The case of two tunnel barriers with $T_{1,n}, T_{2,n} \ll 1$ and for $L_1 = L_2 = L_N$, $r_{N_1} = r_{N_2}$, $R_1 = R_2$, $r_{N_1} \xi_N / L_1 R_1 = 0.0025$ and $\sqrt{2} r_L \xi_S / L R_1 = 0.005$.

Zero-bias anomaly. Let us first analyze the tunneling limit $T_{1,n}, T_{2,n} \ll 1$. In this case at subgap energies the term $g_1^{BTK}(\varepsilon)$ (10) can be neglected and for $E_{1,2} \equiv D_{1,2}/L_{1,2}^2 \ll |\varepsilon| < \Delta$ we obtain

$$g_{11}(\varepsilon) = \frac{\Delta^2}{\Omega^2} \left[\frac{r_{\xi_S}(\varepsilon) + r_{\xi_1}(\varepsilon)}{2R_1^2} + \frac{r_{\xi_S}(\varepsilon)}{2R_1 R_2} e^{-\frac{|x_2 - x_1|}{\xi_S(\varepsilon)}} \right] \quad (16)$$

$$g_{12}(\varepsilon) = \frac{\Omega^2}{2\Delta^2} r_{\xi_S}(\varepsilon) g_{11}(\varepsilon) g_{22}(\varepsilon) e^{-\frac{|x_2 - x_1|}{\xi_S(\varepsilon)}}. \quad (17)$$

Here $r_{\xi_S}(\varepsilon) = r_L \xi_S(\varepsilon)/L$ and $r_{\xi_{1,2}}(\varepsilon) = r_{N_{1,2}} \xi_{1,2}(\varepsilon)/L_{1,2}$ are Drude resistances of the segments of S- and N-metals with respective lengths $\xi_S(\varepsilon) = \sqrt{D_S/2\Omega}$ and $\xi_{1,2}(\varepsilon) = \sqrt{D_{1,2}/|\varepsilon|}$.

At small energies the local spectral conductance diverges as $g_{11}(\varepsilon) \propto 1/\sqrt{\varepsilon}$ which is just well known disorder-induced zero-bias anomaly^{4,5,6}. For the non-local conductance (17) this divergence turns out to be even stronger, $g_{12}(\varepsilon) \propto 1/\varepsilon$, since quantum interference in *both* diffusive normal metals simultaneously enhances non-local electron transport in our system. Thus, we predict a *sharp low energy peak* in the non-local conductance which occurs in the presence of disorder in the N-terminals, see also Fig. 2. Accordingly, the differential conductance $G_{12}(V_2, T) = -\partial I_1 / \partial V_2$ increases as $G_{12} \propto 1/\max(eV_2, T)$ with decreasing voltage and temperature.

Eq. (16) applies down to $\varepsilon \sim E_1$ and for even smaller energies $r_{\xi_1}(\varepsilon)$ should be substituted by $2r_{N_1}$. Then for $r_{N_1}, r_{N_2} \gg r_{\xi_S} \equiv r_{\xi_S}(0)$ we get

$$g_{12}(0) = G_{12}(0, 0) = \frac{r_{\xi_S} r_{N_1} r_{N_2}}{2R_1^2 R_2^2} e^{-|x_2 - x_1|/\xi_S}. \quad (18)$$

We also note that in the case of strongly asymmetric barriers $R_2 \ll R_1$ the dominating contribution to g_{12} scales as $\propto 1/R_1 R_2^3$ rather than $\propto 1/R_1^2 R_2^2$.

Turning to the case of high barrier transmissions $T_{1,n}, T_{2,n} \lesssim 1$ we observe that in this case g_1 is dominated by g_1^{BTK} (10) while other contributions can be neglected. In particular, for fully open barriers at subgap




FIG. 3: (a) Non-local resistance $R_{12}(T)$ (20), normalized by $R_0 = r_{x_1} r_{L-x_2}/r_L$, at different barrier transmissions τ and at $\tau_{Q^*}, \tau_\varphi \rightarrow \infty$. We chose $R_1 = R_2 = \pi/e^2 N_{ch} \tau = \hbar/20e^2$ and set $L = 20\sqrt{2}\xi_S(0)$, $x_1 = 9\sqrt{2}\xi_S(0)$, $x_2 = 11\sqrt{2}\xi_S(0)$, $r_L = 200 \Omega$ and $r_{N_1} = r_{N_2} = 0$. (b) $R_{12}(T)/R_0$ at different $|x_2 - x_1|$. Other parameters are the same as in Fig. 2b.

energies we obtain $g_{1,2} = g_{1,2}^{BTK} = 2/R_{1,2}$ and, hence,

$$g_{12}(\varepsilon) = \frac{\Omega^2}{\Delta^2} \frac{2r_{\xi_S}(\varepsilon)}{R_1 R_2} e^{-|x_2 - x_1|/\xi_S(\varepsilon)}, \quad |\varepsilon| < \Delta. \quad (19)$$

Non-local resistance and charge imbalance peak. Let us now define non-local linear resistance

$$R_{12}(T) = \frac{G_{12}(0, T)}{G_{11}(0, T)G_{22}(0, T) - G_{12}(0, T)G_{21}(0, T)}. \quad (20)$$

Combining this equation with Eq. (14), at $T \ll \Delta$ we arrive at a very simple and universal formula

$$R_{12} = (r_{\xi_S}/2) e^{-|x_2 - x_1|/\xi_S}. \quad (21)$$

It is remarkable that *independently of both barrier and N -terminal parameters* the subgap non-local resistance is set only by the normal state resistance r_{ξ_S} of the superconducting wire segment of length ξ_S and by the distance between the barriers measured in units of ξ_S . At

low T the dependence $R_{1,2} \approx r_0 \exp[-|x_2 - x_1|/\xi_S(0)]$ was observed in experiments¹⁰ with $r_0 \approx 0.56 \Omega$. For the parameters¹⁰ we estimate $r_0 = r_{\xi_S}/2$ in the range of one Ω . A similarly good agreement is found between Eq. (21) and experimental results⁸.

The temperature dependence of $R_{12}(T)$ is depicted in Fig. 3. In the tunneling limit it exhibits a well pronounced peak which originates from the competition between charge imbalance and Andreev reflection^{20,21}. The maximum value of the non-local resistance R_{12} is reached at $T^* \simeq 2\Delta/\ln(R_1 R_2/r_{\xi_S}^2)$ and reads

$$R_{12}(T^*) \approx \frac{\alpha r_{x_1} \sqrt{\frac{8T^*}{\pi\Delta}} \left(1 - \frac{|x_2 - x_1|}{\lambda}\right)}{\left(\sqrt{\frac{r_{\xi_S} + r_{\xi_1}(T^*)}{R_1}} + \sqrt{\frac{r_{\xi_S} + r_{\xi_2}(T^*)}{R_2}}\right)^2}, \quad (22)$$

where $\lambda = \alpha L$ and $\alpha = 1 - r_{x_1}/r_L$. Thus, the peak resistance $R_{12}(T^*)$ decreases linearly with increasing distance $|x_2 - x_1|$ between the barriers. This behavior agrees well with recent observations¹⁰. Furthermore, with the parameters¹⁰ we estimate λ to be of order a micron in agreement with experimental findings. Such values of λ appear lower than typical values of the charge imbalance relaxation length $\Lambda_{Q^*} = \sqrt{D_S \tau_{Q^*}}$. The latter length scale is expected to gain importance only for $\Lambda_{Q^*} < \lambda$.

In summary, we developed a microscopic theory of non-local electron transport in diffusive NSN systems which accounts for non-trivial interplay between crossed Andreev reflection, disorder, quantum interference and non-local charge imbalance. Our results can be directly used for quantitative analysis of future experiments.

This work was supported in part by RFBR grant 09-02-00886. D.S.G. and M.S.K. also acknowledge support respectively from DFG Center for Functional Nanostructures (CFN) and from the Dynasty Foundation.

- ¹ M. Tinkham and J. Clarke, Phys. Rev. Lett. **28**, 1366 (1972); M. Tinkham, Phys. Rev. B **6**, 1747 (1972).
- ² A.F. Andreev, Sov. Phys. JETP **19**, 1228 (1964).
- ³ G.E. Blonder, M. Tinkham, and T.M. Klapwijk, Phys. Rev. B **25**, 4515 (1982).
- ⁴ A.F. Volkov, A.V. Zaitsev, and T.M. Klapwijk, Physica C **210**, 21 (1993).
- ⁵ F.W.J. Hekking and Yu.V. Nazarov, Phys. Rev. Lett. **71**, 1625 (1993); Phys. Rev. B **49**, 6847 (1994).
- ⁶ A.D. Zaikin, Physica B **203**, 255 (1994).
- ⁷ J.M. Byers and M.E. Flatte, Phys. Rev. Lett. **74**, 306 (1995); G. Deutscher and D. Feinberg, Appl. Phys. Lett. **76**, 487 (2000).
- ⁸ D. Beckmann, H.B. Weber, and H. v. Löhneysen, Phys. Rev. Lett. **93**, 197003 (2004); D. Beckmann and H. v. Löhneysen, Appl. Phys. A **89**, 603 (2007).
- ⁹ S. Russo, M. Kroug, T.M. Klapwijk, and A.F. Morpurgo, Phys. Rev. Lett. **95**, 027002 (2005).
- ¹⁰ P. Cadden-Zimansky and V. Chandrasekhar, Phys. Rev.

- Let. **97**, 237003 (2006); P. Cadden-Zimansky, Z. Jiang, and V. Chandrasekhar, New J. Phys. **9**, 116 (2007).
- ¹¹ A. Kleine, A. Baumgartner, J. Trbovic, and C. Schönenberger, arXiv:0812.3553.
- ¹² G. Falci, D. Feinberg, and F.W.J. Hekking, Europhys. Lett. **54**, 255 (2001).
- ¹³ M.S. Kalenkov and A.D. Zaikin, Phys. Rev. B **75**, 172503 (2007); *ibid.* **76**, 224506 (2007).
- ¹⁴ A. Levy Yeyati, F.S. Bergeret, A. Martin-Rodero, and T.M. Klapwijk, Nat. Phys. **3**, 455 (2007).
- ¹⁵ D.S. Golubev and A.D. Zaikin, arXiv:0902.2864.
- ¹⁶ R. Melin, Phys. Rev. B **73**, 174512 (2006).
- ¹⁷ S. Duhot and R. Melin, Eur. Phys. J. B **53**, 257 (2006).
- ¹⁸ A. Brinkman and A.A. Golubov, Phys. Rev. B **74**, 214512 (2006).
- ¹⁹ J.P. Morten, A. Brataas, and W. Belzig, Phys. Rev. B **74**, 214510 (2006).
- ²⁰ D.S. Golubev and A.D. Zaikin, Phys. Rev. B **76**, 184510 (2007).

- ²¹ M.S. Kalenkov and A.D. Zaikin, JETP Lett. **87**, 140 (2008).
- ²² See, e.g., W. Belzig, F. Wilhelm, C. Bruder, G. Schön, and A.D. Zaikin, Superlatt. Microstruct. **25**, 1251 (1999).
- ²³ Yu.V. Nazarov, Superlatt. Microstruct. **25**, 1221 (1999).